



# PRG-2009: Relative Protein Quantification in a Clinical Matrix



R.E. Settlage<sup>1</sup>, A.S. Chien<sup>2</sup>, D.B. Friedman<sup>3</sup>, D.H. Hawke<sup>4</sup>; J. Krijgsveld<sup>5</sup>, K.S. Lilley<sup>6</sup>, M.J. MacCoss<sup>7</sup>, N.E. Sherman<sup>8</sup>, C.W. Turck<sup>9</sup>

<sup>1</sup>Virginia Bioinformatics Institute, Blacksburg VA; <sup>2</sup>Stanford Univ., Stanford CA; <sup>3</sup>Vanderbilt Univ., Nashville TN; <sup>4</sup>Univ. of Texas, MD Anderson Cancer Center, Houston TX;

<sup>5</sup>Utrecht Univ., Utrecht, The Netherlands; <sup>6</sup>Univ. of Cambridge, Cambridge, UK; <sup>7</sup>Univ. of Washington, Seattle WA; <sup>8</sup>Univ. of Virginia, Charlottesville VA; <sup>9</sup>Max Planck Institute of Psychiatry, Munich, Germany

## Proteomics Research Group

## Proteomics Research Group

**Introduction**  
 The Proteomics Research Group (PRG) of the ABRF developed the 2009 study to assess approaches that individual laboratories would use to determine the relative abundance of target proteins in a complex mixture. An increasingly common request for proteomics laboratories is the detection of a specific target protein of interest in a complex mixture. Likewise, most of these requests are also interested in knowing the abundance of the target protein relative to that in a control sample. While this type of analysis has traditionally been addressed using Western blots or other immunobiology assays, recent advances in targeted mass spectrometry-based analyses are beginning to be reported in the literature as an alternative.

For this year's study, four different proteins were spiked into a plasma background matrix at three different levels. Two of these proteins are commonly measured plasma protein biomarkers, and the remaining two had identical primary structure and differed by only a single phosphorylation site. The participants were shipped six samples in total (three samples in blinded duplicate) and asked to report the relative abundances of the four target proteins in the six samples. Results from analysis of the samples and survey responses will be used to assess the different approaches that are used by the proteomics community to determine the relative abundance of a target protein of interest.

**Objectives**  
 The primary goals of this study are to document the breadth of approaches used by the scientific community and to highlight the type of information obtained. Participants were asked to provide the following:

- Relative quantification of four specified proteins in human plasma.
- Information about methods used to analyze the samples.
- Information about the experimental design used for quantification.

**Sample Preparation**  
 Human plasma was obtained from Biomedical Biological Services Inc. (Wilmington, DE). Prostate Specific Antigen (PSA) was obtained from BIOTREND Chemikalien GmbH (Koeln, Germany), chronic gonadotrophin ( $\beta$ -hCG), and glycogen phosphorylase (GP-a/b) were obtained from Sigma Aldrich (St. Louis, MO). Protein stock solutions were prepared at a concentration of 1 mg/mL in 50 mM Tris-HCl, pH 8. Study samples were prepared by spiking appropriate volumes of stock solutions to give target concentrations shown in Table 1 and aliquots were prepared with the volumes shown. The study samples were then lyophilized to dryness and shipped to the study participants for analysis. The participant key is shown in Table 2.

Table	PSA	$\beta$ -hCG	GP-a	GP-b	GP(a+b)	Volume of sample aliquot shipped
	fmol/1L	fmol/1L	fmol/1L	fmol/1L	fmol/1L	$\mu$ L
A and E	1.50 (0.3)	0.75 (0.3)	0.25 (0.1)	1.25 (0.5)	2.00 (1.0)	40.0
B and F	25 (0.5)	12.5 (0.5)	4.0 (0.2)	20.0 (1.0)	32.0 (1.6)	40.0
C and D	250 (10)	125 (5.0)	40 (2.0)	200 (10)	320 (16)	40.0

Table 1: spike concentrations in samples (concentration and normalized)

**Acknowledgements**  
 We would like to thank:  
 Peter Sharratt, University of Cambridge, for amino acid analysis;  
 Giuseppina Maccaroni and Christiane Rewerts, Max Planck Institute of Psychiatry, for preparing test samples;  
 Ms. Karolina Krasinska, Stanford University, for anonymizing the study participants;  
 BiH-Fang Pan, MD Anderson Cancer Center, for performing Western Blots and other tests on initial samples to ensure sample quality;  
 Julie Coleman and Saisha Hill, Vanderbilt University, for sample testing;  
 Kristi Nelson, Virginia Commonwealth University for acquiring MS/MS spectra on the protein standards.


## Target Proteins in the Study.

**Figure 1:** Proteins used in PRG-2009. Please note that all proteins were prepared by commercial vendors and the PRG does not have complete details on proteins. Heterogeneity resulting from truncations, isoforms, post-translational modifications, etc.


**Prostate specific antigen from human seminal fluids (PSA)**  
 may include isoforms 1-6

**human chorionic gonadotropin (recombinant protein from mouse cell line) (a-CG and  $\beta$ -CG)**  
 includes both  $\alpha$  and  $\beta$  isoforms.

**glycogen phosphorylase A/B (from rabbit muscle)**  
 includes both A and B isoforms, B isoforms is phosphorylated






## Participant demographics







## MS instrumentation and methods used

## Lab methods used



## Results and Conclusions



Tally of duplicate calls

Target	sample A	sample B	sample C	sample D	sample E	sample F
1	0.0	0.0	0.0	0.0	0.0	0.0
2	0.0	0.0	0.0	0.0	0.0	0.0
3	0.0	0.0	0.0	0.0	0.0	0.0
4	0.0	0.0	0.0	0.0	0.0	0.0
5	0.0	0.0	0.0	0.0	0.0	0.0
6	0.0	0.0	0.0	0.0	0.0	0.0
7	0.0	0.0	0.0	0.0	0.0	0.0
8	0.0	0.0	0.0	0.0	0.0	0.0
9	0.0	0.0	0.0	0.0	0.0	0.0
10	0.0	0.0	0.0	0.0	0.0	0.0
11	0.0	0.0	0.0	0.0	0.0	0.0
12	0.0	0.0	0.0	0.0	0.0	0.0
13	0.0	0.0	0.0	0.0	0.0	0.0
14	0.0	0.0	0.0	0.0	0.0	0.0
15	0.0	0.0	0.0	0.0	0.0	0.0
16	0.0	0.0	0.0	0.0	0.0	0.0
17	0.0	0.0	0.0	0.0	0.0	0.0
18	0.0	0.0	0.0	0.0	0.0	0.0
19	0.0	0.0	0.0	0.0	0.0	0.0
20	0.0	0.0	0.0	0.0	0.0	0.0
21	0.0	0.0	0.0	0.0	0.0	0.0
22	0.0	0.0	0.0	0.0	0.0	0.0
23	0.0	0.0	0.0	0.0	0.0	0.0
24	0.0	0.0	0.0	0.0	0.0	0.0
25	0.0	0.0	0.0	0.0	0.0	0.0
26	0.0	0.0	0.0	0.0	0.0	0.0
27	0.0	0.0	0.0	0.0	0.0	0.0
28	0.0	0.0	0.0	0.0	0.0	0.0

Relative amount of  $\beta$ -hCG reported by clinical labs

Target	sample A	sample B	sample C	sample D	sample E	sample F
1	0.0	0.0	0.0	0.0	0.0	0.0
2	0.0	0.0	0.0	0.0	0.0	0.0
3	0.0	0.0	0.0	0.0	0.0	0.0
4	0.0	0.0	0.0	0.0	0.0	0.0
5	0.0	0.0	0.0	0.0	0.0	0.0
6	0.0	0.0	0.0	0.0	0.0	0.0
7	0.0	0.0	0.0	0.0	0.0	0.0
8	0.0	0.0	0.0	0.0	0.0	0.0
9	0.0	0.0	0.0	0.0	0.0	0.0
10	0.0	0.0	0.0	0.0	0.0	0.0
11	0.0	0.0	0.0	0.0	0.0	0.0
12	0.0	0.0	0.0	0.0	0.0	0.0
13	0.0	0.0	0.0	0.0	0.0	0.0
14	0.0	0.0	0.0	0.0	0.0	0.0
15	0.0	0.0	0.0	0.0	0.0	0.0
16	0.0	0.0	0.0	0.0	0.0	0.0
17	0.0	0.0	0.0	0.0	0.0	0.0
18	0.0	0.0	0.0	0.0	0.0	0.0
19	0.0	0.0	0.0	0.0	0.0	0.0
20	0.0	0.0	0.0	0.0	0.0	0.0
21	0.0	0.0	0.0	0.0	0.0	0.0
22	0.0	0.0	0.0	0.0	0.0	0.0
23	0.0	0.0	0.0	0.0	0.0	0.0
24	0.0	0.0	0.0	0.0	0.0	0.0
25	0.0	0.0	0.0	0.0	0.0	0.0
26	0.0	0.0	0.0	0.0	0.0	0.0
27	0.0	0.0	0.0	0.0	0.0	0.0
28	0.0	0.0	0.0	0.0	0.0	0.0

Relative amount of PSA reported by clinical labs

Target	sample A	sample B	sample C	sample D	sample E	sample F
1	0.0	0.0	0.0	0.0	0.0	0.0
2	0.0	0.0	0.0	0.0	0.0	0.0
3	0.0	0.0	0.0	0.0	0.0	0.0
4	0.0	0.0	0.0	0.0	0.0	0.0
5	0.0	0.0	0.0	0.0	0.0	0.0
6	0.0	0.0	0.0	0.0	0.0	0.0
7	0.0	0.0	0.0	0.0	0.0	0.0
8	0.0	0.0	0.0	0.0	0.0	0.0
9	0.0	0.0	0.0	0.0	0.0	0.0
10	0.0	0.0	0.0	0.0	0.0	0.0
11	0.0	0.0	0.0	0.0	0.0	0.0
12	0.0	0.0	0.0	0.0	0.0	0.0
13	0.0	0.0	0.0	0.0	0.0	0.0
14	0.0	0.0	0.0	0.0	0.0	0.0
15	0.0	0.0	0.0	0.0	0.0	0.0
16	0.0	0.0	0.0	0.0	0.0	0.0
17	0.0	0.0	0.0	0.0	0.0	0.0
18	0.0	0.0	0.0	0.0	0.0	0.0
19	0.0	0.0	0.0	0.0	0.0	0.0
20	0.0	0.0	0.0	0.0	0.0	0.0
21	0.0	0.0	0.0	0.0	0.0	0.0
22	0.0	0.0	0.0	0.0	0.0	0.0
23	0.0	0.0	0.0	0.0	0.0	0.0
24	0.0	0.0	0.0	0.0	0.0	0.0
25	0.0	0.0	0.0	0.0	0.0	0.0
26	0.0	0.0	0.0	0.0	0.0	0.0
27	0.0	0.0	0.0	0.0	0.0	0.0
28	0.0	0.0	0.0	0.0	0.0	0.0

Normalized ratio GP-(b/a)

Target	sample A	sample B	sample C	sample D	sample E	sample F
1	0.0	0.0	0.0	0.0	0.0	0.0
2	0.0	0.0	0.0	0.0	0.0	0.0
3	0.0	0.0	0.0	0.0	0.0	0.0
4	0.0	0.0	0.0	0.0	0.0	0.0
5	0.0	0.0	0.0	0.0	0.0	0.0
6	0.0	0.0	0.0	0.0	0.0	0.0
7	0.0	0.0	0.0	0.0	0.0	0.0
8	0.0	0.0	0.0	0.0	0.0	0.0
9	0.0	0.0	0.0	0.0	0.0	0.0
10	0.0	0.0	0.0	0.0	0.0	0.0
11	0.0	0.0	0.0	0.0	0.0	0.0
12	0.0	0.0	0.0	0.0	0.0	0.0
13	0.0	0.0	0.0	0.0	0.0	0.0
14	0.0	0.0	0.0	0.0	0.0	0.0
15	0.0	0.0	0.0	0.0	0.0	0.0
16	0.0	0.0	0.0	0.0	0.0	0.0
17	0.0	0.0	0.0	0.0	0.0	0.0
18	0.0	0.0	0.0	0.0	0.0	0.0
19	0.0	0.0	0.0	0.0	0.0	0.0
20	0.0	0.0	0.0	0.0	0.0	0.0
21	0.0	0.0	0.0	0.0	0.0	0.0
22	0.0	0.0	0.0	0.0	0.0	0.0
23	0.0	0.0	0.0	0.0	0.0	0.0
24	0.0	0.0	0.0	0.0	0.0	0.0
25	0.0	0.0	0.0	0.0	0.0	0.0
26	0.0	0.0	0.0	0.0	0.0	0.0
27	0.0	0.0	0.0	0.0	0.0	0.0
28	0.0	0.0	0.0	0.0	0.0	0.0

**Discussion/Conclusions**  
 Study samples were prepared as shown in Table 1 and distributed to 67 participants. The study participants were asked to determine the relative abundance of 4 proteins spiked into the samples and report their methods and results.

As a control, two sample sets were analyzed for PSA and beta HCG by clinical laboratories using standard ELISA techniques. As shown above, the results of the clinical laboratories were largely in agreement with our theoretical values calculated using quantitation values from amino acid analysis. Both of the clinical laboratories did note some difficulty in analyzing PSA levels in samples A and E. In addition, PSA levels in samples B and F appeared to be below the detection limit of CL-2.

While most successful respondents found depletion of the most abundant proteins helpful, success was achieved using all instrument types interfaced with chromatographic separation with and without depletion. All successful participants used enzymatic digestion (trypsin). Successful MS scanning methods included SR/MS/MS, peptide precursor/product signal extracted from full scan MS/MS, and MS<sup>1</sup> methods. Most participants did not find sample amount limiting.

Several participants noted some difficulty with the lyophilized state of the samples. It should be noted that samples were lyophilized prior to shipping for the sole purpose of reducing the cost of the study.

This year's study was rated as difficult to very difficult by the participants with the average confidence level of the participants ability to perform the analysis remaining at confident to very-confident before and after the study. When asked whether they would repeat the study given an opportunity, we received a resounding 'yes, it was fun' with many requests for additional time to perform the study.

Table 2: study participant key

Participant	Target 1	Target 2	Target 3	Target 4
1	1	1	1	1
2	1	1	1	1
3	1	1	1	1
4	1	1	1	1
5	1	1	1	1
6	1	1	1	1
7	1	1	1	1
8	1	1	1	1
9	1	1	1	1
10	1	1	1	1
11	1	1	1	1
12	1	1	1	1
13	1	1	1	1
14	1	1	1	1
15	1	1	1	1
16	1	1	1	1
17	1	1	1	1
18	1	1	1	1
19	1	1	1	1
20	1	1	1	1
21	1	1	1	1
22	1	1	1	1
23	1	1	1	1
24	1	1	1	1
25	1	1	1	1
26	1	1	1	1
27	1	1	1	1
28	1	1	1	1
29	1	1	1	1
30	1	1	1	1
31	1	1	1	1
32	1	1	1	1
33	1	1	1	1
34	1	1	1	1
35	1	1	1	1
36	1	1	1	1
37	1	1	1	1
38	1	1	1	1